

**Table 1 Composition of the Bulgarian grain legume collections**

Genus	Total	Long term conservation	Middle term conservation	Working ex situ collection
<i>Pisum</i>	2420 / <i>120</i>	610 / <i>200</i>	900 / 250	1000 / <i>120</i>
<i>Lupinus</i>	296 / <i>12</i>	70	204	10
<i>Lathyrus</i>	337 / <i>17</i>	270	100	40 / <i>17</i>
<i>Phaseolus</i>	1763/ <i>120</i>	777	1151	138 / <i>120</i>
<i>Cicer</i>	348	234	56	4
<i>Lens</i>	532	385	237	94
<i>Vicia sp.</i>	1780 / <i>200</i>	1150	534	124 / <i>54</i>
<i>Soja hispida</i>	700	692	-	7
<i>Vicia faba</i>	692/ <i>10</i>	462	220	5 / <i>10</i>
<i>Vigna</i>	270	117	123	88
<i>Arachis</i>	685	317 / 43	368	368

• *progress number*

**Table2 Ex-situ collection of *Vicia sp.* with Bulgarian origin**

Species	Varieties	Landraces	Wild relatives.	Total
<i>V. sativa</i> complex	48	162	118	328
<i>Vicia villosa</i>	1	3	20	24
<i>Vicia narbonensis</i>			16	16
<i>Vicia pannonica</i>			21	21
<i>Vicia ervillia</i>	1	2	1	4
Other			169	169
<b>Total</b>	<b>50</b>	<b>167</b>	<b>345</b>	<b>562</b>

fig 1




Fig.2




fig.3

